

表1 2006年の評価法使用頻度（全体）

3編以上の論文に使用された評価法を示す。

| 評価法 | 論文数 |
|--|-----|
| Functional independence measure, FIM | 50 |
| Mini-mental state examination, MMSE | 30 |
| Medical outcomes study short form-36 health survey, SF-36 | 27 |
| Barthel index, BI, バーセル指数 | 25 |
| Ashworth scale-modified, MAS | 17 |
| Berg balance scale, BBS | 13 |
| Fugl-Meyer assessment | 14 |
| 6 minute walk test, 6MWT | 13 |
| Timed up and go, TUG | 13 |
| Gross motor function measure, GMFM | 7 |
| National Institutes of Health stroke scale, NIHSS | 8 |
| Activities-specific balance confidence scale, ABC | 7 |
| American Spinal Injury Association impairment scale, ASIA impairment scale | 7 |
| Geriatric depression scale, GDS | 6 |
| Ashworth scale | 6 |
| Brunnstrom stage | 6 |
| Community integration questionnaire, CIQ | 6 |
| Hospital anxiety and depression scale, HAD | 7 |
| Motricity index, MI | 6 |
| Glasgow coma scale, GCS | 5 |
| Sickness impact profile, SIP | 5 |
| Trail making test, TMT | 5 |
| Action research arm test, ARAT | 5 |
| Borg scale | 4 |
| Fatigue severity scale, FSS | 4 |
| Functional assessment measure, FAM | 4 |
| Modified Rankin scale | 4 |
| Oswestry disability index | 4 |
| Rivermead mobility index, RMI | 3 |
| Trunk control test, TCT | 4 |
| Beck depression inventory, BDI | 3 |
| Disability rating scale, DRS | 3 |
| Dizziness handicap inventory, DHI | 3 |
| Expanded disability status scale, EDSS | 3 |
| Frenchay activities index, FAI | 3 |
| Functional Ambulation Categories, FAC | 3 |
| International classification of functioning, disability and health, ICF | 4 |
| Katz ADL index | 3 |
| Motor assessment scale, MAS | 3 |
| Pediatric evaluation of disability inventory, PEDI | 3 |
| Rivermead motor assessment | 3 |
| Gross Motor Function Classification System, GMFCS | 3 |