

表1 評価法使用頻度と論文のエビデンスレベル (2005年)

3篇以上の論文に使用された評価法を示す。

評価法	合計	エビデンスレベル					
		Ia	Ib	IIa	IIb	III	IV
10(6) meter walk test (10MWT)	6		2	2	1	1	
2(6) minute walking test (2MWT)	6				2	4	
Action research arm test, ARAT	3		1		1	1	
American Spinal Injury Association impairment scale, ASIA impairment scale	22		1	1	6	12	2
Ashworth scale	10		1	2		6	1
Ashworth scale-modified, MAS	16		3	4	4	5	
Barthel index, BI, バーセル指数	30		2	1	10	13	4
Beck depression inventory, BDI	5				3	2	
Berg balance scale (BBS)	6			2		4	
Borg scale	5		1	1	1	2	
Brunnstrom stage	5		1		2	1	1
Center for epidemiological studies-depression scale, CES-D	5		1	1		3	
Chedoke-McMaster Stroke Assessment (CMSA)	3		1		1	1	
Comprehensive severity index (CSI)	7				7		
Disability rating scale (DRS)	7		2		4	1	
EuroQol	3	1	1	1			
Fear avoidance beliefs questionnaire (FABQ)	3		1		1	1	
Fugl-Meyer assessment	9	1	1	2	4	1	
Functional autonomy measurement system, SMAF	3					3	
Functional independence measure for children, WeeFIM	3					3	
Functional independence measure, FIM	78	1	5	7	27	31	7
Geriatric depression scale, GDS	4			1	2	1	
Glasgow coma scale, GCS	15		2		6	7	
Glasgow outcome scale, GOS	4		1	1	1	1	
Gross motor function classification system	5			1	1	3	
Hamilton anxiety rating scale (HARS)	3		1		2		
Health assessment questionnaire	3				2	1	
Hoehn-Yahr stage/score	3				3		
Hospital anxiety and depression scale	3				2	1	
International classification of diseases, 9th revision, ICD-9	10				8	2	
Japan coma scale, JCS	4					3	1
Kohs 立方体検査	3						3
Manual muscle testing, MMT, 徒手筋力検査	10				1	2	7
Medical outcomes study short form-12 health survey, SF-12	4				2	1	1
Medical outcomes study short form-36 health survey, SF-36	24	2	4	0	4	14	
Mini-mental state examination, MMSE	24			2	6	12	4
Motor activity log, MAL	4		1	2		1	
Neck Disability Index (NDI)				2	1		
Nottingham health profile, NHP	3		1			2	
Numeric rating scale, NRS	3		1		1	1	
Oswestry disability index	6	1	1	1	2	1	
Pediatric Evaluation of Disability Inventory (PEDI)	4			1	2	1	
Raven colored matrices	3				1		2
Rivermead Post Concussion Symptoms Questionnaire (RPQ)	3			1		2	
Roland and Morris disability questionnaire, RMDQ	6	1	3		1	1	
Short Portable Mental Status Questionnaire (SPMSQ)	3				1	2	
Sickness impact profile, SIP	4	2				2	
Stroke impairment assessment set, SIAS	7			1	2	3	1
Timed up & go (TUG) test	6		1	3	2		
Trail making test, TMT	7			1	1	1	4
Visual Analog Scale (VAS)	17		3	2	5	6	1
Wechsler Adult Intelligence Scale-Revised, WAIS-R, ウェクスラー成人知能検査	7			1	2	1	3
Wisconsin card sorting test, WCST	3				1		2
簡易上肢機能テスト, STEF	3				1		2
長谷川式簡易痴呆スケール, HDS	6				2	2	2