

表1 2006年の評価法使用頻度（全体）

3編以上の論文に使用された評価法を示す。

評価法	論文数
Functional independence measure, FIM	50
Mini-mental state examination, MMSE	30
Medical outcomes study short form-36 health survey, SF-36	27
Barthel index, BI, バーセル指数	25
Ashworth scale-modified, MAS	17
Berg balance scale, BBS	13
Fugl-Meyer assessment	14
6 minute walk test, 6MWT	13
Timed up and go, TUG	13
Gross motor function measure, GMFM	7
National Institutes of Health stroke scale, NIHSS	8
Activities-specific balance confidence scale, ABC	7
American Spinal Injury Association impairment scale, ASIA impairment scale	7
Geriatric depression scale, GDS	6
Ashworth scale	6
Brunnstrom stage	6
Community integration questionnaire, CIQ	6
Hospital anxiety and depression scale, HAD	7
Motricity index, MI	6
Glasgow coma scale, GCS	5
Sickness impact profile, SIP	5
Trail making test, TMT	5
Action research arm test, ARAT	5
Borg scale	4
Fatigue severity scale, FSS	4
Functional assessment measure, FAM	4
Modified Rankin scale	4
Oswestry disability index	4
Rivermead mobility index, RMI	3
Trunk control test, TCT	4
Beck depression inventory, BDI	3
Disability rating scale, DRS	3
Dizziness handicap inventory, DHI	3
Expanded disability status scale, EDSS	3
Frenchay activities index, FAI	3
Functional Ambulation Categories, FAC	3
International classification of functioning, disability and health, ICF	4
Katz ADL index	3
Motor assessment scale, MAS	3
Pediatric evaluation of disability inventory, PEDI	3
Rivermead motor assessment	3
Gross Motor Function Classification System, GMFCS	3