

表8 1998年以来の全データでの使用頻度  
(1998～2000年, 2003～2006年)

| 評価法  | 論文数 |
|--|-----|
| Functional independence measure, FIM                                       | 379 |
| Barthel index, BI, バーセル指数  | 153 |
| Mini-mental state examination, MMSE  | 118 |
| American Spinal Injury Association impairment scale, ASIA impairment scale | 117 |
| Medical outcomes study short form-36 health survey, SF-36                  | 113 |
| Ashworth scale-modified, MAS   | 78  |
| Glasgow coma scale, GCS  | 77  |
| Brunnstrom stage   | 61  |
| Fugl-Meyer assessment  | 54  |
| Ashworth scale   | 48  |
| Wechsler Adult Intelligence Scale-Revised, WAIS-R, ウェクスラー成人知能検査            | 46  |
| Manual muscle testing, MMT, 徒手筋力検査   | 45  |
| Berg balance scale (BBS)   | 45  |
| Beck depression inventory, BDI   | 41  |
| Sickness impact profile, SIP   | 38  |
| Timed up & go (TUG) test   | 36  |
| Disability rating scale (DRS)  | 32  |
| 6-minute walking test (6MWT)   | 31  |
| Trail making test, TMT   | 30  |
| International classification of diseases, 9th revision, ICD-9              | 26  |
| 長谷川式簡易痴呆スケール, HDS  | 25  |
| Community integration questionnaire, CIQ                                   | 25  |
| Stroke impairment assessment set, SIAS                                     | 25  |
| Hospital anxiety and depression scale, HADS                                | 23  |
| Geriatric depression scale, GDS  | 21  |
| Glasgow outcome scale, GOS   | 20  |
| Nottingham health profile, NHP   | 20  |
| Roland and Morris disability questionnaire, RMDQ                           | 20  |